

# The impact of COVID-19 prevention measures on surgical wound infection rates post-cardiac surgery

Aladdin Bashir, Matthew Holmes, Nebumathew Suresh, Pedram Panahi, Sameh Atta, Hannah T. Perkins, Clinton Lloyd, Sanjay Asopa

University Hospitals Plymouth NHS Trust, Plymouth, United Kingdom

## Abstract

The COVID-19 pandemic had a huge impact on medical services. Several measures have been implemented to reduce the risk of viral transmission. In this paper, we assessed the impact of these measures on surgical wound infection rates in post-cardiac surgery

Correspondence: Aladdin Bashir, University Hospitals Plymouth, NHS Trust, Plymouth, United Kingdom.  
E-mail: aladdin.bashir@nhs.net

Key words: cardiac surgery, wound infection, COVID-19, infection control, postoperative complications.

Contributions: all authors have contributed significantly and agree with the content of the manuscript. All authors have read and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

Conflict of interest: the authors declare no conflict of interest.

Ethics approval and consent to participate: the study protocol was approved by the Audit, Assurance & Effectiveness Team of the University Hospitals of Plymouth (protocol no. CA\_2021-22-120 dated 17/09/2021).

Patient consent for publication: not applicable.

Funding: none.

Availability of data and materials: the data used to support the findings of this study are available from the corresponding author upon request.

Received: 12 April 2023.  
Accepted: 10 June 2023.  
Early view: 12 October 2023.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2023  
Licensee PAGEPress, Italy  
Monaldi Archives for Chest Disease 2024; 94:2604  
doi: 10.4081/monaldi.2023.2604

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial International License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

patients. Hypothesis testing was used to compare post-cardiac operation infection rates between the year prior to the COVID-19 pandemic being declared and the first 13 months of the pandemic. The infection rates in 969 patients with operations between 01/03/2019 and 29/02/2020 were compared to those of 925 patients with cardiac surgery between 01/03/2020 and 31/03/2021. Infection rates for various operative urgencies and infection types were analyzed. To compare infection rates, a two-tailed pooled z-test using the difference in infection proportions was performed. A 5% significance level was used, and only categories with at least 10 patients in both the pre-COVID and COVID populations were tested. For leg infections, only operations involving coronary artery bypass grafting were included. To ensure that any differences in outcomes were not due to differences in patient demographics resulting in unequal operative risks, Euroscore II values, a measure of cardiac operative risk, were compared between the pre-COVID and post-COVID cohorts. The Mann-Whitney U-test was used to determine whether the distributions of Euroscore II values were likely to be drawn from the same population. A significance level of 5% was used. A total of 1901 patients (932 during the COVID-19 pandemic) were included in this study. There was a significant reduction in post-operative infections for all patients undergoing cardiac surgery, from 4.3% of patients before COVID-19 to 1.5% during the pandemic. During the pandemic, fewer elective and more urgent operations were performed. This study suggests a significant role for iatrogenic causes in wound infections before the pandemic. The implementation of COVID-19 prevention measures by healthcare providers can reduce surgical infection rates. As COVID-19-related restrictions have been eased, we suggest maintaining them for healthcare providers to reduce the incidence of surgical wound infections.

## Introduction

Surgical site infection (SSI) post-cardiac surgery can have a detrimental impact on patient morbidity and mortality [1]. They are also associated with an increased length of hospital stay, long-term antibiotic use, further surgical intervention, and increased costs [1-3]. SSI in patients' post-cardiac surgery can include sternal wound infections (SWI) and leg wound infections in patients requiring coronary artery bypass grafting (CABG). Fortunately, the incidence of SWI is relatively low in comparison to other surgical wounds, with an incidence ranging from 0.5% to 6.0% [2,4]. There is an extensive list of modifiable and non-modifiable risk factors which include: poor glucose control, *Staphylococcus aureus* skin colonization, smoking, inadequate skin preparation, hypo- or hyperthermia, and hypoxia [1,5]. The World Health Organization has released guidelines that endorse the use of skin barriers, skin decontamination, and intraoperative homeothermy, in addition to hand hygiene

measures for the prevention of SSI [6]. Many of these measures were more rigorously implemented during the COVID-19 pandemic in 2020/2021.

The COVID-19 pandemic had a sweeping impact on all aspects of society, most notably healthcare. In an attempt to curb the transmission of the virus in healthcare settings and therefore protect both patients and staff members, a variety of measures were implemented. Hospitals prohibited visitors from visiting their relatives, canceled routine work, and crucially enforced scrupulous hygiene measures. This included more frequent hand washing, fewer physical interactions between healthcare workers and patients, and greater use of personal protective equipment (PPE). These measures represent the basic level of infection control precautions that should be always applied during the care of all patients. Infection prevention and control during health care when COVID-19 is suspected or confirmed.

Hand hygiene is one of the most effective precautionary measures to prevent the spread of the virus [7]. The rationale and the correct use of PPE played a leading role in slowing down the virus spread. The proper training of the staff, the appropriate selection of different types of masks according to the situation, and the high compliance of the staff and the community establish hygienic habits in the hospital setting [8,9].

Despite the effects of the pandemic, our center was still performing cardiac surgery, mostly for patients requiring urgent or emergency operations. It was noted locally that the rates of SSIs appeared to be lower during the COVID-19 pandemic. Therefore, this study aims to compare the rates of SSI during the COVID-19 pandemic when intensified hygiene measures were being implemented with those rates observed before the pandemic.

## Materials and Methods

Hypothesis testing was used to compare post-cardiac operation infection rates between the year before the COVID-19 pandemic was declared and the first 13 months of the pandemic. The infection rates in 969 patients with operations between 01/03/2019 and 29/02/2020 were compared to those of 925 patients with cardiac surgery between 01/03/2020 and 31/03/2021. Infection rates for various operative urgencies and infection types were analyzed. To compare infection rates, a two-tailed pooled z-test using the difference in infection proportions was performed. A 5% significance level was used, and only categories with at least 10 patients in both the pre-COVID and COVID populations were tested. For leg infections, only operations involving CABG were included.

To ensure that any differences in outcomes were not due to differences in patient demographics resulting in unequal operative risks, Euroscore II values (Figure 1), a measure of cardiac operative risk, were compared between the pre-COVID and post-COVID cohorts. The Mann-Whitney U-test was used to determine whether the distributions of Euroscore II values were likely to be drawn from the same population. This test is analogous to the *t*-test but does not make the assumption that the underlying distribution is normal. In this case, the distributions were frequently long-tailed. A significance level of 5% was used. For leg infections, only operations involving CABG were included.

## Results

There was a significant reduction in post-operative infections for all patients undergoing cardiac surgery, from 4.3% of patients before

COVID-19 to 1.5% during the pandemic (Figures 2 and 3). The proportion of patients developing a post-operative infection is similar or reduced when comparing pre- and during-pandemic operations for all urgency levels and infection categories (where sufficient data was available for testing). Notably, during the pandemic, fewer elective and more urgent operations were performed (Figure 4).

## Discussion and Conclusions

SSI in cardiac surgery can be categorized by location and depth. Both graft site infections and SWI can be classified as either superficial or deep. Superficial SWI involves the skin, subcutaneous tissue, and pectoralis fascia, whereas deep SWI (DSWI), or mediastinitis, involves any tissue deep into the subcutaneous tissue [1]. This can include any of the great vessels, trachea, esophagus, and the heart itself. The incidence of DSWI is reported to be 0.2-3% compared to 2-6% for superficial infections [10,11]. Although relatively uncommon, they have a significant effect on morbidity, mortality, and the duration of hospital stays [1,2,3]. One-year mortality rates have been reported to be around 10% for patients with DSWI compared to controls [12], with hospital stays reported to be four times longer (32 days) [13]. DSWI is predominantly caused by *Staphylococcus* species such as coagulase-negative *Staphylococci* and *Staphylococcus aureus* [14], although other species such as *Escherichia coli*, *Pseudomonas aeruginosa*, and *Klebsiella* are also commonly cultured [15]. There are a number of perioperative measures that can be implemented to reduce the risk of developing such infections. Patients are routinely screened for nasal carriers of *S. aureus* to identify those that need eradication therapy and appropriate prophylactic antibiotics. Patients should also be optimized in relation to glycemic control, nutrition, and smoking [14].


Diabetes mellitus and obesity are both well-established risk factors for developing DSWI. Multiple studies have demonstrated diabetes as a significant risk factor, and those diabetics with a pre-operative blood glucose concentration  $\geq 11.1$  mmol/L are as much as ten times more likely to develop DSWI [16-18]. Obesity increases the odds of DSWI by up to 2.6 times [19]. The same systematic review found no significant relationship between smoking and sternal wound infection; however, only a small number of studies were included. Numerous papers have found it to be an independent risk factor for developing other SSI, and smaller studies have found it to be a risk factor for DSWI [20,21]. Routine nasal swabs for *Staphylococcus* colonization and routine intranasal mupirocin administration in combination with chlorhexidine gluconate bathing in the absence of nasal cultures or nasal cultures positive have been shown to significantly decrease the incidence of DSWI following cardiac surgery [22].

During the pandemic, additional precautions were implemented within all hospital settings in the UK to minimize the risk of spreading COVID-19 (Table 1). These predominantly came in the form of additional PPE (including disposable gloves and aprons) for all clinical staff during patient interactions. The implementation of stricter handwashing policies and reduced skin-to-skin contact (including post-operative physical examinations) was also adopted. Furthermore, face masks were mandatory for all hospital staff in all clinical areas, and visitor numbers and the duration of visits were significantly lower.

There is an abundance of literature focusing on pre-operative and operative techniques to minimize the risks of general SSI and SWI [7,8,14-16,22-24]. However, fewer studies focus on the impact of COVID-19 prevention measures on SSI rates [25-29]. Our results are consistent with the findings by Hussain *et al.* [25], who noticed

a decrease in the incidence of sternal wound infection during the pandemic. It has been demonstrated that the consumption of PPE and products during the COVID-19 pandemic has led to a decrease in SSI in patients after cesarean delivery [26] as well as after spinal surgery [27]. Hand hygiene is considered the cornerstone of the prevention of surgical wound infections [7,23]. It has been shown that hand hygiene's quality and frequency have significantly improved

during the COVID-19 pandemic, which led to a reduction in hospital infections [27,30]. The use of PPE was an essential addition to daily practice during the COVID-19 pandemic. Non-medical (fabric) masks were used by all workers working indoors or in close proximity to clients and co-workers, as per the World Health Organization guidelines [31]. Despite the wide use of disposable surgical masks, their efficacy in wound infection prevention is unclear, and data are



**Figure 1.** Euroscore II distributions: distributions statistically similar enough to have come from the same parent distribution for the following urgency categories: urgent, emergency, elective & urgent, emergency and salvage and all.

limited [32]. Nevertheless, the significant reduction in wound infection rates could be related to the routine use of PPE during the COVID-19 pandemic [27].

While some of the COVID-19 prevention measures have been

eased (physical distancing, family visits), this paper highlights the benefits of keeping others (rigorous hand hygiene, PPE in clinical areas, frequent cleaning, and disinfection of environmental surfaces) in surgical departments.

Operative Urgency	Infection Type	Number of Operations Pre Covid	Number of Operations Post Covid	Infection Proportion Pre Covid	Infection Proportion Post Covid	p-value for Differences in Infection Proportion	Euroscore II Welsch t-test p-value	Significant Difference in Euroscore II Distributions?
Elective	Superficial Chest	631	388	0.00792393	0.005154639	0.603277483	0.059680687	Yes
Elective	Deep Chest	631	388	0.020602219	0.00257732	0.016387107	0.059680687	Yes
Elective	Mediastinitis	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Elective	Superficial Leg	333	206	0.018018018	0.009708738	0.438213586	0.059680687	Yes
Elective	Deep Leg	333	206	0.006006006	0	0.265115735	0.059680687	Yes
Elective	Any Chest	631	388	0.026941363	0.005154639	0.012542325	0.059680687	Yes
Elective	Any Leg	333	206	0.024024024	0.009708738	0.231383737	0.059680687	Yes
Elective	Any	631	388	0.041204437	0.010309278	0.004611082	0.059680687	Yes
Urgent	Superficial Chest	281	461	0.007117438	0.00867679	0.818047595	0.321319391	No
Urgent	Deep Chest	281	461	0.021352313	0.006507592	0.073155198	0.321319391	No
Urgent	Mediastinitis	281	461	0.007117438	0.002169197	0.302858274	0.321319391	No
Urgent	Superficial Leg	216	359	0.018518519	0	0.009669885	0.321319391	No
Urgent	Deep Leg	216	359	0	0.002785515	0.437542284	0.321319391	No
Urgent	Any Chest	281	461	0.03202847	0.017353579	0.19499401	0.321319391	No
Urgent	Any Leg	216	359	0.018518519	0.002785515	0.049082571	0.321319391	No
Urgent	Any	281	461	0.049822064	0.019522777	0.020888847	0.321319391	No
Emergency	Superficial Chest	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency	Deep Chest	48	68	0	0.014705882	0.398773883	0.712882564	No
Emergency	Mediastinitis	48	68	0.020833333	0	0.231930076	0.712882564	No
Emergency	Superficial Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency	Deep Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency	Any Chest	48	68	0.020833333	0.014705882	0.80282027	0.712882564	No
Emergency	Any Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency	Any	48	68	0.020833333	0.014705882	0.80282027	0.712882564	No
Salvage	Superficial Chest	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Salvage	Deep Chest	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Salvage	Mediastinitis	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Salvage	Superficial Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Salvage	Deep Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Salvage	Any Chest	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Salvage	Any Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Salvage	Any	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Elective & Urgent	Superficial Chest	912	849	0.007675439	0.007067138	0.881548039	0.999377851	No
Elective & Urgent	Deep Chest	912	849	0.020833333	0.004711425	0.002905654	0.999377851	No
Elective & Urgent	Mediastinitis	912	849	0.002192982	0.001177856	0.605745527	0.999377851	No
Elective & Urgent	Superficial Leg	549	565	0.018214936	0.003539823	0.017682064	0.999377851	No
Elective & Urgent	Deep Leg	549	565	0.003642987	0.001769912	0.546440923	0.999377851	No
Elective & Urgent	Any Chest	912	849	0.028508772	0.011778563	0.013173027	0.999377851	No
Elective & Urgent	Any Leg	549	565	0.021857923	0.005309735	0.016582033	0.999377851	No
Elective & Urgent	Any	912	849	0.043859649	0.015312132	0.000458974	0.999377851	No
Emergency & Salvage	Superficial Chest	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency & Salvage	Deep Chest	57	83	0	0.012048193	0.405590953	0.57224968	No
Emergency & Salvage	Mediastinitis	57	83	0.01754386	0	0.225880344	0.57224968	No
Emergency & Salvage	Superficial Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency & Salvage	Deep Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency & Salvage	Any Chest	57	83	0.01754386	0.012048193	0.787760853	0.57224968	No
Emergency & Salvage	Any Leg	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Emergency & Salvage	Any	57	83	0.01754386	0.012048193	0.787760853	0.57224968	No
All	Superficial Chest	969	932	0.007223942	0.006437768	0.835287461	0.272571438	No
All	Deep Chest	969	932	0.019607843	0.005364807	0.005427223	0.272571438	No
All	Mediastinitis	969	932	0.003095975	0.001072961	0.335914936	0.272571438	No
All	Superficial Leg	568	588	0.017605634	0.003401361	0.017214955	0.272571438	No
All	Deep Leg	568	588	0.003521127	0.00170068	0.543056721	0.272571438	No
All	Any Chest	969	932	0.027863777	0.011802575	0.012378876	0.272571438	No
All	Any Leg	568	588	0.021126761	0.005102041	0.016092068	0.272571438	No
All	Any	969	932	0.042311662	0.015021459	0.000387124	0.272571438	No

Figure 2. Cardiac surgical patients stratified by operative urgency and wound infection type. NA, not available.



Operative Urgency	Infection Type	Number of Operations Pre Covid	Number of Operations Post Covid	Infection Proportion Pre Covid	Infection Proportion Post Covid	p-value for Differences in Infection	Euroscore II Welsch t-test p-value	Significant Difference in Euroscore II
Urgent	Superficial Leg	216	359	0.018518519	0	0.009669885	0.321319391	No
Urgent	Any Leg	216	359	0.018518519	0.002785515	0.049082571	0.321319391	No
Urgent	Any	281	461	0.049822064	0.019522777	0.020888847	0.321319391	No
Elective & Urgent	Deep Chest	912	849	0.020833333	0.004711425	0.002905654	0.999377851	No
Elective & Urgent	Superficial Leg	549	565	0.018214936	0.003539823	0.017682064	0.999377851	No
Elective & Urgent	Any Chest	912	849	0.028508772	0.011778563	0.013173027	0.999377851	No
Elective & Urgent	Any Leg	549	565	0.021857923	0.005309735	0.016582033	0.999377851	No
Elective & Urgent	Any	912	849	0.043859649	0.015312132	0.000458974	0.999377851	No
All	Deep Chest	969	932	0.019607843	0.005364807	0.005427223	0.272571438	No
All	Superficial Leg	568	588	0.017605634	0.003401361	0.017214955	0.272571438	No
All	Any Chest	969	932	0.027863777	0.011802575	0.012378876	0.272571438	No
All	Any Leg	568	588	0.021126761	0.005102041	0.016092068	0.272571438	No
All	Any	969	932	0.042311662	0.015021459	0.000387124	0.272571438	No

Figure 3. Statistically significant differences in infection rates.




Figure 4. Operative urgency.

Table 1. COVID-19 prevention measures applied in our trust during the pandemic.

Measure	Pre COVID	During COVID
Physical distancing	Not applied	Applied at least 6 inches
Hand hygiene	After patient physical contact and after touching any soiled equipment	Mandatory after dealing with any patients up to patient files
Cleaning and disinfection of environmental surfaces	Routine decontamination and sterilization of surfaces and equipment	More frequent decontamination of surfaces, walls, keyboards, personal equipment including phones and luggage of the patients and the healthcare professionals
Personal protective equipment	Only in designated clinical areas (theatres, intensive care)	In all clinical and non-clinical areas with an upgrade to higher protective equipment (FFP3) when dealing with COVID positive patients or in COVID wards
Family visiting	Relaxed visiting policy with no restriction on time or number of visitors	It varied during the COVID period and ranged from non-visiting policy, moving to limited number and timing with COVID negative proof

## References

1. Sharif M, Wong CHM, Harky A. Sternal wound infections, risk factors and management-how far are we? A literature review. *Heart Lung Circ* 2019;28:835-43.
2. Schiraldi L, Jabbour G, Centofanti P, et al. Deep sternal wound infections: Evidence for prevention, treatment, and reconstructive surgery. *Arch Plast Surg* 2019;46:291-302.
3. Cayci C, Russo M, Cheema F, et al. Risk analysis of deep sternal wound infections and their impact on long-term survival: a propensity analysis. *Ann Plast Surg* 2008;61:294-301.
4. Hever P, Singh P, Eiben I, et al. The management of deep sternal wound infection: literature review and reconstructive algorithm. *JPRAS Open* 2021;28:77-89.
5. Jayakumar S, Khoynezhad A, Jahangiri M. Surgical site infections in cardiac surgery. *Crit Care Clin* 2020;36:581-92.
6. World Health Organization. Global guidelines for the prevention of surgical site infection. Available from: <https://apps.who.int/iris/handle/10665/277399>.
7. World Health Organization. Patient safety. WHO guidelines on hand hygiene in health care: first global patient safety challenge clean care is safer care. Available from: [https://apps.who.int/iris/bitstream/handle/10665/44102/9789241597906\\_eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/44102/9789241597906_eng.pdf).
8. World Health Organization. Rational use of personal protective equipment (PPE) for coronavirus disease (COVID-19): interim guidance, 19 March 2020. Available from: <https://apps.who.int/iris/handle/10665/331498>.
9. Honda H, Iwata K. Personal protective equipment and improving compliance among healthcare workers in high-risk settings. *Curr Opin Infect Dis* 2016;29:400-6.
10. Crabtree TD, Codd JE, Fraser VJ, et al. Multivariate analysis of risk factors for deep and superficial sternal infection after coronary artery bypass grafting at a tertiary care medical center. *Semin Thorac Cardiovasc Surg* 2004;16:53-61.
11. Ridderstolpe L, Gill H, Granfeldt H, et al. Superficial and deep sternal wound complications: incidence, risk factors and mortality. *Eur J Cardiothorac Surg* 2001;20:1168-75.
12. Sears ED, Wu L, Waljee JF, et al. The impact of deep sternal wound infection on mortality and resource utilization: a population-based study. *World J Surg* 2016;40:2673-80.
13. Graf K, Ott E, Vonberg RP, et al. Economic aspects of deep sternal wound infections. *Eur J Cardiothorac Surg* 2010;37:893-6.
14. Phoon PHY, Hwang NC. Deep sternal wound infection: diagnosis, treatment and prevention. *J Cardiothorac Vasc Anesth* 2020;34:1602-13.
15. Pradeep A, Rangasamy J, Varma PK. Recent developments in controlling sternal wound infection after cardiac surgery and measures to enhance sternal healing. *Med Res Rev* 2021;41:709-24.
16. Trick WE, Scheckler WE, Tokars JI, et al. Modifiable risk factors associated with deep sternal site infection after coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2000;119:108-14.
17. Martin ET, Kaye KS, Knott C, et al. Diabetes and risk of surgical site infection: a systematic review and meta-analysis. *Infect Control Hosp Epidemiol* 2016;37:88-99.
18. Furu M, Kong PK, Moorthy PSK, et al. Risk factors for sternal wound infection after coronary artery bypass grafting in patients with and without diabetes. *Int Heart J* 2022;63:426-32.
19. Balachandran S, Lee A, Denehy L, et al. Risk factors for sternal complications after cardiac operations: a systematic review. *Ann Thorac Surg* 2016;102:2109-17.
20. Colombier S, Kessler U, Ferrari E, et al. Influence of deep sternal wound infection on long-term survival after cardiac surgery. *Med Sci Monit* 2013;19:668-73.
21. Olsen MA, Lock-Buckley P, Hopkins D, et al. The risk factors for deep and superficial chest surgical-site infections after coronary artery bypass graft surgery are different. *J Thorac Cardiovasc Surg* 2002;124:136-45.
22. Bode LGM, Kluytmans JAJW, Wertheim HFL, et al. Preventing surgical-site infections in nasal carriers of *Staphylococcus aureus*. *N Engl J Med* 2010;362:9-17.
23. Allegranzi B, Pittet D. Role of hand hygiene in healthcare-associated infection prevention. *J Hosp Infect* 2009;73:305-15.
24. Assadian O, Golling M, Krüger CM, et al. Surgical site infections: guidance for elective surgery during the SARS-CoV-2 pandemic - international recommendations and clinical experience. *J Hosp Infect* 2021;111:189-99.
25. Hussain A, Ike DI, Durand-Hill M, Ibrahim S, Roberts N. Sternal wound infections during the COVID-19 pandemic: an unexpected benefit. *Asian Cardiovasc Thorac Ann* 2021;29:376-80.
26. Antonello VS, Dallé J, Antonello ICF, et al. Surgical site infection after cesarean delivery in times of COVID-19. *Rev Bras Ginecol Obstet* 2021;43:374-6.
27. Perna A, Maruccia F, Gorgoglione FL, et al. Increased frequency of hand hygiene and other infection prevention practices correlates with reduced surgical wound infection rates in spinal surgery during the COVID-19 pandemic. *J Clin Med* 2022;11:7528.
28. Sybert M, Oakley CT, Christensen T, et al. Impact of COVID-19 protocols on primary and revision total hip arthroplasty. *J Arthroplasty* 2022;37:2193-8.
29. Lazar HL. Commentary: compliance with the American Association for Thoracic Surgery guidelines will prevent sternal wound infections and minimize postoperative complications in cardiac surgery patients during the COVID-19 pandemic. *J Thorac Cardiovasc Surg* 2020;160:e44-8.
30. Roshan R, Feroz AS, Rafique Z, Virani N. Rigorous hand hygiene practices among health care workers reduce hospital-associated infections during the COVID-19 pandemic. *J Prim Care Community Health* 2020;11:2150132720943331.
31. World Health Organization. Preventing and mitigating COVID-19 at work: policy brief, 19 May 2021. Available from: <https://apps.who.int/iris/handle/10665/341328>.
32. Vincent M, Edwards P. Disposable surgical face masks for preventing surgical wound infection in clean surgery. *Cochrane Database Syst Rev* 2016;4:CD002929.