

# Assessment of left ventricular function in subjects with chronic obstructive pulmonary disease

Murali Mupparapu,<sup>1</sup> Rajesh Venkataram,<sup>2\*</sup> Nandakishore Baikunje,<sup>2</sup> Ivor D'Sa,<sup>3</sup> Chandramouli Mandya Thimmaiah<sup>2</sup>

<sup>1</sup>Department of Nephrology, Basaveshwara Medical College and Hospital, Chitradurga, Karnataka; <sup>2</sup>Department of Pulmonary Medicine, K S Hegde Medical Academy, Mangalore, Karnataka; <sup>3</sup>Department of General Medicine, K S Hegde Medical Academy, Mangalore, Karnataka, India

\*Present address: Department of Pulmonary Medicine, Yenepoya Medical College Mangalore, Karnataka, India

Correspondence: Chandramouli Mandya Thimmaiah, Department of Pulmonary Medicine, K S Hegde Medical Academy, Mangalore 575018, Karnataka, India.  
E-mail: mouli.aims@gmail.com

Key words: COPD, LV diastolic dysfunction, LV systolic dysfunction, echocardiography, E/A.

Contributions: MM, conceptualization, data curation, formal analysis, methodology, original draft; NB, RV, ID, formal analysis, investigation, methodology, visualization, data curation; CMT, formal analysis, validation, visualization, review, and editing. All the authors read and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

Conflict of interest: the authors declare no potential conflict of interest.

Ethics approval and consent to participate: the study was approved by the Institutional Ethics Committee of KS Hegde Medical Academy which is a constituent institution of Nitte (Deemed to be University) approval ID: INST.EC/EC/146/2014-15 dated 13.10.2014. Informed consent was obtained from all individual participants included in the study.

Informed consent: written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article. The manuscript does not contain any individual person's data in any form.

Patient consent for publication: obtained.

Availability of data and materials: the data used to support the findings of this study are available from the corresponding author upon request.

Funding: none.

Received: 5 June 2023.

Accepted: 29 September 2023.

Early view: 16 October 2023.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Monaldi Archives for Chest Disease 2024; 94:2654

doi: 10.4081/monaldi.2023.2654

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial International License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## Abstract

Chronic obstructive pulmonary disease (COPD) is the world's third leading cause of death, posing a significant public health challenge. Heart failure is an important milestone in the natural history of this disease's ever-decreasing course. Cor pulmonale alone is sometimes insufficient to explain the scenario, as many heart failures are refractory even after standard treatment. A possible explanation is the involvement of the left ventricle (LV). However, there is currently no consensus on a therapeutic approach to LV dysfunction in COPD. Some previous studies were conducted on COPD patients regardless of the presence or absence of other factors influencing LV function. This study attempted to rule out all known confounding factors/comorbidities that influence LV function. We discovered that LV diastolic dysfunction was common in subjects at all stages of COPD. We believe that all COPD patients, regardless of stage, should have screening echocardiography.

## Introduction

Chronic obstructive pulmonary disease (COPD), though preventable and treatable, is currently being ranked the third leading cause of death worldwide [1]. Globally, the reported prevalent COPD cases were 212.3 million with 3.3 million deaths and 74.4 million disability-adjusted life-years in 2019 [2]. In India, the estimated prevalence is 7.4%. It is responsible for more than 50 lakh deaths yearly, which is 4 times higher than that in the Western world [3]. The socio-economic implications of such a disease cannot be overlooked.

On the morbidity front, exercise limitation is the hallmark of this disease which cannot be fully explained by lung function parameters alone. Deranged chest wall mechanics, dysfunction of the diaphragm and skeletal muscles of the limbs, and cardiac abnormalities like cor pulmonale are some contributors to exercise limitation [4]. Among these, cor pulmonale heralds the onset of the rapid downhill course of the disease. However, despite prompt treatment, heart failure frequently remains refractory, and left ventricular involvement could be the possible explanation. It is well known that left ventricle (LV) filling dynamics are influenced by right ventricular (RV) loading conditions. Abnormal patterns of LV diastolic filling have been reported in the setting of increased RV pressure and/or volume. This has also been noted in the absence of other disorders affecting LV function [5]. Still, no guidelines have thoroughly addressed this issue. Hence, the present study was conducted to examine LV dysfunction in subjects with COPD in our tertiary care referral hospital.

## Materials and Methods

This was a prospective study conducted at our tertiary care hospital, from October 2014 to June 2018. It involved consenting subjects diagnosed with COPD based on post-bronchodilator ratio of the forced expiratory volume in the first one second to the forced vital capacity (FEV1/FVC) <0.7 on spirometry as per the Global Initiative for Chronic Obstructive Lung Disease (GOLD) criteria. Subjects who were diagnosed with acute exacerbation of COPD, who had co-existing cardiac disease, who had systemic hypertension, diabetes mellitus, dyslipidemia, who were unable to perform spirometry, who had a history of alcohol consumption >2 drinks/day in females and >3 drinks/day in males (known to cause cardiomyopathy) and who were not consenting, were excluded. Demographic, clinical, hematological, spirometry, electrocardiogram (ECG), echocardiography, and chest X-ray data were gathered. COPD was staged as below.

Post-bronchodilator FEV1/FVC<0.7 with i) post bronchodilator FEV1≥80% predicted -GOLD stage 1 (mild); ii) 50%≤FEV1<80% predicted - GOLD stage 2 (moderate); iii) 30%≤FEV1<50% predicted - GOLD stage 3 (severe); iv) FEV1<30% - GOLD stage 4 (very severe).

The cardiologist performed 2-dimensional transthoracic Doppler echocardiography.

E<sub>max</sub> (E) was defined as early maximal trans-mitral flow velocity and A<sub>max</sub> (A) was defined as late peak atrial systolic velocity. E/A was used as an indicator of LV diastolic function. LV ejection fraction was used as an indicator of LV systolic dysfunction. Left ventricular end-systolic diameter and left ventricular end-diastolic diameter were used as indicators of ventricular remodeling.

### Grading of diastolic dysfunction

Grade 1 (mild dysfunction): E/A ratio of less than 0.75 m/s.

Grade 2 (moderate dysfunction): E/A ratio of 1-1.5 m/s and normal deceleration time (160-240 ms).

Grade 3 and 4 (severe irreversible dysfunction): E/A ratio higher than 2 m/s.

### Statistical analysis

Data collected was entered in SPSS 20 (IBM, Armonk, NY, USA). A descriptive analysis was done. The chi-square test, analysis of variance, and Pearson's correlation coefficient calculation were done. A p-value <0.05 was considered significant. Approval from the institutional ethics committee was obtained.

## Results

During the study period, 410 consecutive subjects with COPD were identified. Overall, 38 nonconsenting subjects, 42 subjects who could not perform spirometry, and 261 subjects with co-existing cardiac disease or other confounding factors were excluded. Thus, 69 eligible subjects were studied. Among them, 49 subjects were male and 20 were female. The predominant age distribution was 50-70 years. Most of the subjects belonged to GOLD stage 2 and stage 3. Among 69 subjects, 9 had normal ventricular function, and 60 had evidence of ventricular dysfunction. Out of 60 subjects, 50 (72.46%) had evidence of LV diastolic dysfunction (LVDD), 5 had LV systolic dysfunction, and the remaining 5 subjects had cor pulmonale (Table 1).

The incidence of LVDD (E/A) with increasing age was not significant (p>0.05). However, E<sub>max</sub> as defined by early maximal transmitral flow velocity alone correlated well with increasing age (Table 2). Symptoms such as cough with sputum production, breathlessness, smoking status, and duration of symptoms did not correlate with LV dysfunction (Table 3). However, recent worsening of breathlessness, dizziness, and syncope were the only symptoms that had a significant statistical correlation with LVDD (Table 4).

## Discussion

LVDD in subjects with COPD may be due to LV pre-load and afterload abnormalities. Chest wall hyperinflation secondary to emphysema may affect LV diastolic function by impairing LV fill-

**Table 1.** Demographic details, Global Initiative for Chronic Obstructive Lung Disease staging, and echocardiogram findings.

Particulars	n	%
Age (in years)		
41-50	9	13.5
51-60	25	36.23
61-70	25	36.23
>70	10	14.49
Sex		
Male	49	71
Female	20	29
GOLD staging		
Grade 1	11	15.94
Grade 2	27	39.13
Grade 3	22	31.88
Grade 4	9	13.03
ECHO findings		
Normal	9	13.06
LV diastolic dysfunction	50	72.46
LV systolic dysfunction	5	7.24
Cor pulmonale	5	7.24

GOLD, Global Initiative for Chronic Obstructive Lung Disease; ECHO, echocardiogram; LV, left ventricle.

**Table 2.** Correlation between age and E<sub>max</sub>.

		E <sub>max</sub>	Age
E <sub>max</sub>	Pearson correlation	1	-0.300*
	Sig. (2-tailed)		0.020
	N	60	60
Age	Pearson correlation	-0.300*	1
	Sig. (2-tailed)	0.020	
	N	60	60

**Table 3.** Correlation with duration of symptoms with presence of left ventricle dysfunction.

		E/A	Symptoms
E/A	Pearson correlation	1	0.086
	Sig. (2-tailed)		0.513
	N	60	60
Symptoms	Pearson Correlation	0.086	1
	Sig. (2-tailed)	0.513	
	N	60	60

ing and pre-load [6-8]. Intrinsic positive end-expiratory pressure is common in subjects with COPD, which is known to limit venous return. Its severity is generally proportional to the severity of hyperinflation and airway obstruction [9]. Reduction of aortic diastolic blood pressure may reduce coronary perfusion, leading to sub-endocardial ischemia. Both effects lead to myocardial fibrosis with impaired myocardial relaxation [10]. Thus, these subjects will likely have ventricular and arterial stiffening with impaired systolic and diastolic function in the absence of overt cardiovascular disease [11]. In this context, our findings suggest a potential mechanism underscoring the excess risk of coronary heart disease and congestive heart failure in COPD and the possibility of a relationship between airway obstruction, arterial stiffness, and cardiovascular disease in the general population [12,13]. Also, subjects with COPD experience chronic hypoxemia, which might result in abnormalities of myocardial relaxation as a consequence of intracellular calcium transport disturbances [14]. Systemic inflammation may provide a biological link between the two entities, as COPD has been regarded as a systemic disease with various extrapulmonary manifestations. Tumor necrosis factor  $\alpha$ , and interleukins 6 and 18 have been implicated as the key inflammatory mediators [15].

In our study, we found that a significant proportion (72.46%) of subjects with COPD had LVDD. Our findings are consistent with those of Rawy and Fathalla [16], who also found LVDD to be the predominant form of LV dysfunction, being detected in 73.3% of the subjects with COPD. Vizza *et al.* have observed LV systolic dysfunction in less than 5% of subjects with COPD [17]. Malerba *et al.* have observed LVDD in 65% of subjects with COPD [18]. Steinberg *et al.* have found LVDD in 78% of subjects with COPD [19]. A few similar studies have been summarized in Table 5 [20-24]. A possible explanation for this observation might be that impaired myocardial relaxation and fibrosis (discussed above) lead to diastolic dysfunction early in the course of the disease before the subject develops systolic dysfunction.

We did not find any correlation between LV dysfunction and the presence of respiratory symptoms such as cough with expectoration, chest pain, and breathlessness, or with a duration of these symptoms (Table 3). We found a statistically significant correlation only in subjects reporting recent worsening breathlessness and complaints of the cardiovascular system such as syncope and dizziness (Table 4). It is to be noted that dizziness or syncope are not common symptoms associated with COPD. However, Gaude *et al.* [16] found a significant correlation between the presence of LV dysfunction and the duration of symptoms, though, it is unclear whether echocardiography was performed during an acute exacerbation. It may be noted that myocardial strain present in subjects who have acute breathlessness may be a factor leading to false positive evidence of LVDD [25]. In our study, an echocardiogram (ECHO) was not done dur-

ing acute episodes to avoid this confounding effect. Since tachycardia also influences diastolic function, ECHO was delayed till the heart rate was less than 100 bpm.

Age is a proven risk factor for LV dysfunction. We found that age did not significantly correlate with E/A ( $p=0.27$ ), which is the indicator of LVDD. Thus, we could rule out age's influence on LV function. However, Emax alone correlated well with increasing age (Table 2). We found that with progression in stages of COPD, there was no change in E/A (Table 6). This implies that there was no proper matching between the presence of LVDD and the stage of COPD. Further, pulmonary hypertension is known to influence the left ventricular dynamics indirectly through its effects on the interventricular septum. Notably, pulmonary hypertension was not present in any of the 50 subjects with LVDD in our study. Funk *et al.* [26], and Boussuges *et al.* [4] have reported similar observations albeit in a lesser proportion of subjects. All the above observations point to the presence of LVDD in the absence of other well-known factors commonly associated with it.

It is well established that pulmonary hypertension starts to appear in the clinical picture as the individual moves to higher grades of COPD. And hence, ECHO is not generally performed until then unless there are other medical indications. In this background, our observations suggest that performing ECHO in COPD subjects irrespective of the GOLD stage may be worth doing for timely detection of LVDD.

We did not find any significant correlation between LV dys-

**Table 4.** Correlations of different variables with left ventricular diastolic dysfunction.

Variable	p
Age	0.270
Symptoms of respiratory system	0.280
Symptoms of cardiovascular system	
Chest discomfort	0.549
Chest pain	0.849
Pedal edema	0.549
Syncope and dizziness	0.042
Breathlessness (recently worsening)	0.050
Smoking	0.393
Tuberculosis in the past	0.177
Duration of symptoms	0.603
GOLD staging	
Stage 1	0.177
Stage 2	0.171
Stage 3	0.115
Stage 4	0.470

GOLD, Global Initiative for Chronic Obstructive Lung Disease.

**Table 5.** Studies with observed proportions of subjects having different forms of left ventricle dysfunction.

Authors	Number of subjects with COPD (n)	Subjects with LV systolic dysfunction (%)	Subjects with LV diastolic dysfunction (%)
Steele <i>et al.</i> [20]	120	21	60
Berger <i>et al.</i> [21]	27	10	72
Mac Nee <i>et al.</i> [22]	36	14	65
Zema <i>et al.</i> [23]	37	36	68
McCullough <i>et al.</i> [24]	417	46	78

COPD, chronic obstructive pulmonary disease; LV, left ventricle.

**Table 6.** Analysis of variance of left ventricle dysfunction with the grading of chronic obstructive pulmonary disease.

ANOVA		Sum of squares	Df	Mean Square	F	Sig.
Emax	Between groups	0.007	3	0.002	0.110	0.954
	Within groups	1.200	56	0.021		
	Total	1.207	59			
Amax	Between groups	0.043	3	0.014	0.226	0.878
	Within groups	3.521	56	0.063		
	Total	3.563	59			
E/A	Between groups	0.057	3	0.019	0.400	0.04
	Within groups	2.661	56	0.048		
	Total	2.718	59			
LVESD	Between groups	111.890	3	37.297	0.457	0.713
	Within groups	4571.093	56	81.627		
	Total	4682.983	59			
LVEDD	Between groups	542.498	3	180.833	3.503	0.021
	Within groups	2890.902	56	51.623		
	Total					

ANOVA, analysis of variance; E/A, Emax/Amax; LVESD, left ventricular end-systolic diameter; LVEDD, left ventricular end-diastolic diameter.

function and risk factors predisposing the patients to COPD such as smoking, and a history of tuberculosis (Table 4).

Even though all patients underwent chest X-rays and ECG, these were assessed by a single expert only. Hence, we did not correlate ECHO findings with chest X-ray or ECG.

Even though our study has limitations, such as being a single-centered, cross-sectional study with a small study population, we have tried to eliminate all the potential confounding factors, such as systemic hypertension, and type 2 diabetes mellitus which are well-known comorbid conditions associated with LV dysfunction apart from age. We followed up with three subjects who were diagnosed with COPD by spirometry and the presence of LVDD on echocardiography and further underwent thread mill test and coronary angiography. These tests were normal, essentially ruling out the cardiac cause for LVDD. However, the number of cases who underwent invasive procedures was too small to conclude. In our tertiary care hospital, where we have more than 750 COPD subjects under follow-up, 69 cases were selected after excluding all risk factors that have an independent correlation with LVDD. Hence, the findings in our study are significant enough to prove the association of COPD with LVDD, if not causation. Early diagnosis and timely intervention in patients with COPD reduce the risk of patients progressing to refractory cardiac failure. Hence, all subjects with COPD should preferably undergo echocardiography at the earliest for evaluation of left ventricular function.

## Conclusions

LVDD is present in a significant proportion of subjects with COPD irrespective of the stage. Useful pointers to LVDD could be symptoms such as recent worsening of breathlessness, dizziness, and syncope. It is worth screening all COPD subjects for LVDD using transthoracic echocardiography.

## References

1. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020;396:1204-22. Erratum in: *Lancet* 2020;396:1562.
2. Safiri S, Carson-Chahhoud K, Noori M, et al. Burden of chronic obstructive pulmonary disease and its attributable risk factors in 204 countries and territories, 1990-2019: results from the Global Burden of Disease Study 2019. *BMJ* 2022;378:e069679.
3. Jindal SK. COPD: The unrecognized epidemic in India. *J Assoc Physicians India* 2012;60:14-6.
4. Boussuges A, Pinet C, Molenat F, et al. Left atrial and ventricular filling in chronic obstructive pulmonary disease. An echocardiographic and Doppler study. *Am J Respir Crit Care Med* 2000;162:670-5.
5. Nootens M, Wolfkiel CJ, Chomka EV, Rich S. Understanding right and left ventricular systolic function and interactions at rest and with exercise in primary pulmonary hypertension. *Am J Cardiol* 1995;75:374-7.
6. Redfield MM, Jacobsen SJ, Borlaug BA, et al. Age- and gender-related ventricular-vascular stiffening: a community-based study. *Circulation* 2005;112:2254-62.
7. Nagueh SF, Middleton KJ, Kopelen HA, et al. Doppler tissue imaging: a noninvasive technique for evaluation of left ventricular relaxation and estimation of filling pressures. *J Am Coll Cardiol* 1997;30:1527-33.
8. Kass DA. Ventricular arterial stiffening: integrating the pathophysiology. *Hypertension* 2005;46:185-93.
9. Aldrich TK, Hendler JM, Vizioli LD, et al. Intrinsic positive end-expiratory pressure in ambulatory patients with airways obstruction. *Am Rev Respir Dis* 1993;147:845-9.
10. Laurent S, Cockcroft J, Van Bortel L, et al. Expert consensus document on arterial stiffness: methodological issues and clinical applications. *Eur Heart J* 2006;27:2588-605.
11. Kawaguchi M, Hay I, Fetis B, Kass DA. Combined ventricular systolic and arterial stiffening in patients with heart failure and preserved ejection fraction: implications for systolic and diastolic reserve limitations. *Circulation* 2003;107:714-20. Erratum in: *Circulation* 2020;141:e809.
12. Curkendall SM, DeLuise C, Jones JK, et al. Cardiovascular disease in patients with chronic obstructive pulmonary disease, Saskatchewan Canada cardiovascular disease in COPD patients. *Ann Epidemiol* 2006;16:63-70.
13. Mattace-Raso FU, van der Cammen TJ, Hofman A, et al. Arterial stiffness and risk of coronary heart disease and stroke: the Rotterdam Study. *Circulation* 2006;113:657-63.

14. Nishimura RA, Appleton CP, Redfield MM, et al. Noninvasive doppler echocardiographic evaluation of left ventricular filling pressures in patients with cardiomyopathies: a simultaneous Doppler echocardiographic and cardiac catheterization study. *J Am Coll Cardiol* 1996;28:1226-33.
15. Zvezdin B, Milutinov S, Kojicic M, et al. A postmortem analysis of major causes of early death in patients hospitalized with COPD exacerbation. *Chest* 2009;136:376-80.
16. Rawy AM, Fathalla D. Left ventricular diastolic dysfunction in patients with chronic obstructive pulmonary disease (COPD), prevalence and association with disease severity: Using tissue Doppler study. *Egypt J Chest Dis Tubercul* 2015;64:785-92.
17. Vizza CD, Lynch JP, Ochoa LL, et al. Right and left ventricular dysfunction in patients with severe pulmonary disease. *Chest* 1998;113:576-83.
18. Malerba M, Ragnoli B, Salameh M, et al. Sub-clinical left ventricular diastolic dysfunction in early stage of chronic obstructive pulmonary disease. *J Biol Regul Homeost Agents* 2011;25:443-51.
19. Steinberg BA, Zhao X, Heidenreich PA, et al. Trends in patients hospitalized with heart failure and preserved left ventricular ejection fraction: prevalence, therapies and outcomes. *Circulation* 2012;126:65-75.
20. Steele P, Ellis JH, Van Dyke D, et al. Left ventricular ejection fraction in severe chronic obstructive airways disease. *Am J Med* 1975;59:21-8.
21. Berger HJ, Matthay RA, Loke J, et al. Assessment of cardiac performance with quantitative radionuclide angiocardiology: right ventricular ejection fraction with reference to findings in chronic obstructive pulmonary disease. *Am J Cardiol* 1978;41:897-905.
22. MacNee W, Xue QF, Hannan WJ, et al. Assessment by radionuclide angiography of right and left ventricular function in chronic bronchitis and emphysema. *Thorax* 1983;38:494-500.
23. Zema MJ, Masters AP, Margouloff D. Dyspnea: the heart or the lungs? Differentiation at bedside by use of the simple Valsalva maneuver. *Chest* 1984;85:59-64.
24. McCullough PA, Hollander JE, Nowak RM, et al. Uncovering heart failure in patients with a history of pulmonary disease: rationale for the early use of B-type natriuretic peptide in the emergency department. *Acad Emerg Med* 2003;10:198-204.
25. Ozben B, Eryuksel E, Tanrikulu AM, et al. Acute exacerbation impairs right ventricular function in COPD patients. *Hellenic J Cardiol* 2015;56:324-31.
26. Funk GC, Lang I, Schenk P, et al. Left ventricular diastolic dysfunction in patients with COPD in the presence and absence of elevated pulmonary arterial pressure. *Chest* 2008;133:1354-9.