



Monaldi Archives for Chest Disease

eISSN 2532-5264

<https://www.monaldi-archives.org/>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community. These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Monaldi Archives for Chest Disease** is, therefore, e-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

All legal disclaimers applicable to the journal apply to this production process as well.

Monaldi Arch Chest Dis 2024 [Online ahead of print]

To cite this Article:

Amador AF, Amorim S, Proença T, et al. **Screening of cardiac allograft vasculopathy in heart transplant patients with coronary computed tomography angiography.** *Monaldi Arch Chest Dis* doi: 10.4081/monaldi.2024.2890

 ©The Author(s), 2024
Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.



Screening of cardiac allograft vasculopathy in heart transplant patients with coronary computed tomography angiography

Ana Filipa Amador,¹ Sandra Amorim,¹ Tânia Proença,¹ Mariana Vasconcelos,¹
Marta Tavares Da Silva,¹ João Rebelo,² André Carvalho,²
José Pinheiro-Torres,³ Paulo Pinho,³ Rui Rodrigues¹

¹Department of Cardiology, São João Hospital and University Center, Porto; ²Department of Radiology, São João Hospital and University Center, Porto; ³Department of Cardiothoracic Surgery, São João Hospital and University Center, Porto, Portugal

Correspondence: Ana Filipa Polido Amador, Department of Cardiology, São João Hospital and University Center, Alameda Hernani Monteiro, Porto, 4200-319, Portugal

Tel.: +351933176428.

E-mail: a.filipa.amador@gmail.com

Contributions: AFA, SA, were responsible for the design of the work, acquisition, analysis, and interpretation of the data, as well as drafting the paper; TP, designed the CT protocol and contributed to the paper draft; MV, was responsible for the design of the work; MV, MTDS, JR, AC, performed the CT and revised of the paper; JPT, PP, RAR, reviewed the paper critically for important intellectual content. All authors finally approved the submitted version and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Conflict of interest: nothing to declare.

Ethics approval and consent to participate: the study was performed according to the recommendations of the Helsinki Declaration and patients' consent was acquired. Observational studies are not mandatory to be proposed to the local ethical committee.

Patient consent for publication: acquired.

Funding: none.

Availability of data and materials: raw data were generated at Hospital Universitário de São João, Porto. Derived data supporting the findings of this study are available from the corresponding author on request. Data was collected anonymously.

Abstract

Although coronary angiography (CA) is the gold standard for coronary allograft vasculopathy (CAV) screening, non-invasive modalities have arisen as potential alternatives, such as coronary computed tomography angiography (CCTA). CCTA also quantifies plaque burden, which may influence medical treatment.

From January 2021 to April 2022, we prospectively included heart transplant recipients who performed CCTA as a first-line method for CAV detection in a single center. Clinical, CCTA, and CA data were collected. 38 patients were included, 60.5% men, aged 58 ± 14 years. The most frequent cause of transplantation was dilated cardiomyopathy (42.1%), and the median graft duration was 10 years [interquartile range (IQR) 9]. The median left ventricle ejection fraction was 61.5% (IQR 6). The median calcium score was 17 (IQR 231) and 32 patients (84.2%) proceeded to CCTA: 7, 24, and 1 patients had a graded CAV of 0, 1, and 2, respectively. Most patients (37.5%) had both calcified and non-calcified plaques, and the median number of affected segments was 2 (IQR 3). The remaining six patients had extensive coronary calcification, so CA was performed: 4 had CAV1, 1 had CAV2, and 1 had CAV3.

During follow-up ($12,2 \pm 4,2$ months), there were neither deaths nor acute coronary syndromes. After CCTA, therapeutic changes occurred in about 10 (26.3%) of patients, mainly related to anti-lipid intensification; such changes were more frequent in patients with diabetes after heart transplant. In this cohort, CCTA led to therapeutic changes in about one-quarter of patients; more studies are needed to assess how CCT may guide therapy according to plaque burden.

Key words: screening, heart transplant, coronary allograft vasculopathy, coronary computed tomography angiography.

Introduction

Cardiac allograft vasculopathy (CAV) is the leading cause of long-term graft dysfunction of the heart transplant (HT) patients [1]. It is an accelerated fibro-proliferative coronary disease whose pathological hallmark is intimal thickening, with progressive luminal narrowing, potentially leading to ischemia, heart failure, arrhythmias and death [2]. According to the 2019 International Heart Lung Transplant Society (ISHLT) report, CAV incidence increases over time, developing in about 30% of HT patients at 5 years and almost 50% at 10 years [3]. ISHLT

grading system for angiographic CAV includes the categories of CAV 0 (non-significant), CAV 1 (mild), CAV 2 (moderate), and CAV 3 (severe) disease, which are designated based on the severity of angiographic stenosis as well as on the presence of allograft dysfunction [1]. HT patients need to be regularly monitored for CAV as it is usually symptomatically silent or atypical due to the heart denervation. At the time of this study, according to latest 2010 ISHLT guidelines, coronary angiography (CA) and/or IVUS was the gold standard for screening of CAV and should be performed annually or biannually (class I, LOE C) [4]. However, its invasive nature and inability to thoroughly examine the distal coronary vessels present is a current limitation. According to these guidelines, coronary computed tomography angiography (CCTA) shows promise in the evaluation of CAV in HT recipients, although higher resting heart rates in these patients limit the technical quality of this study (Class IIb, LOE C) [4]. However CCTA has had new technological advancements, providing high-resolution anatomical images of the lumen, but also on the wall of the coronary arteries. Therefore, CCTA has the potential to be a non-invasive alternative to invasive coronary angiography for the diagnosis of CAV. Besides its role in evaluating CAV, CCTA also quantifies plaque burden, which may influence medical treatment.

In the present article, we describe our experience during the first months of implementation in clinical practice of CCTA as the first-line test for CAV detection, and seek to find how CCT findings impact subsequent medical approach.

Materials and Methods

This prospective observational study included HT recipients who performed a CCTA for routine monitoring of CAV in a tertiary care centre, over a 16-month period from January 2021 to April 2022. If calcium score <400, a CCTA was completed; if it was ≥ 400, a CA was performed as an alternative.

Patients with previous diagnose of CAV2 and 3 were excluded (maintained monitorization with CA). Clinical characteristics including patient's medical history, CAV classification, and medication use were obtained from medical records of the most recent outpatient visit. HT recipient patients with resting high heart rate (>100/min), typical of cardiac denervation, were medicated with 5 mg twice a day ivabradine within the 3 days before the exam and/or had an additional dose of verapamil or metoprolol, to achieve compatible heart rate for obtaining gating images.

Regarding CCTA protocol, a dual-source system was used - SOMATON Definition Flash CT scanner, by Siemens, of 2x128 slices. Either prospective ECG-gated high-pitch acquisition (if heart rate was <70 bpm) or retrospective ECG-gated acquisition (if heart rate was ≥ 70 bpm) were used. For contrast injection, we used contrast agents with high iodine concentrations of

370 mg/ml; time monitoring was made at ascending aorta and with bolus tracking. A biphasic injection was performed, with 50-80 ml (1 ml/Kg/weight) contrast media of 5 ml/s and 40-50 ml saline chaser of 5 ml/s. A representative image of CCTA is depicted at Figure 1.

Results

A total of 38 patients were included, with 23 (60,5%) men and mean age of 58 ± 14 years old. Table 1 lists some clinical characteristics. The main cause of transplantation was familial dilated cardiomyopathy (42,1%) and the median graft duration was 10 years (interquartile range – IQR 9). Only one patient had impaired left ventricle ejection fraction (of 45%) and all had preserved right ventricle systolic function. Most patients had at least one cardiovascular risk factor (89,5%) and took anti-platelet or anti-thrombotic drugs (97,4%). A total of 25 (65,8%) patients had history of a previous rejection: 7 (18,4%) patients had at least one episode of antibody mediated rejection and 24 (63,2%) had at least one episode of cellular rejection; 4 patients had previous cytomegalovirus (CMV) infection.

Regarding immunosuppression, 32 (84,2%) patients received prednisolone and 36 (94%) took calcineurin inhibitors; 18 (47,4%) patients were treated with mycophenolate mofetil/mycophenolic acid and 3 (7,9%) patients with mTOR inhibitors.

At CCTA preparation, three patients with stage 4 chronic kidney disease (CKD) were hospitalized for contrast nephropathy prophylaxis; due to resting high heart rate, 35 (81,6%) received verapamil 72h previous to exam. Additionally, immediately before the exam, 24 (63,2%) patients required verapamil (17 intravenous (iv) and 7 oral) and one patient (2,6%) needed metoprolol iv.

After performing CT calcium score, 32 (84,2%) patients proceed to CCTA – see Table 2 for findings of CCTA. The remaining 6 patients had extensive coronary calcification (400) and a CA was performed in replacement: 4 had CAV1, 1 had CAV2 and 1 had CAV3.

After CCTA, 4 (10,5%) patients needed additional ischemia testing: one that had CAV3, both the patients with CAV2 and one patient with CAV1 but suboptimal quality of CCTA images. All four patients underwent myocardial perfusion scintigraphy (MPS), no significant ischemia was detected for any patient.

During the follow up of $12,2 \pm 4,2$ months, there were no deaths, no percutaneous revascularization, no acute coronary syndrome, no ventricular arrhythmias, or stroke. Three patients had de novo rejection - 1 humoral, 1 cellular and 1 both; the patient with both subtypes of rejection evolved from CAV1 to CAV3, was submitted to coronary artery bypass graft surgery and underwent re-transplantation later.

Overall, there was an association between previous antibody mediated rejection and CAV1/2/3 ($p=0.005$). No association was found between previous cellular rejection and CAV1/2/3 ($p=0.216$) nor previous overall rejection and CAV1/2/3 ($p=0.166$).

Findings in CCTA lead to therapeutic changes in 10 (26,3%) of patients – all ten patients had lipid drugs intensification, with increased statin dose in one patient, switch from low to high intensity statin in 5 patients and addition of ezetimibe in 4 patients. Moreover, anti-thrombotic therapy was changed from platelet anti-aggregation to warfarin due to intra-cardiac thrombus in 3 patients; one patient with CAV1, noncalcified plaques and 21 years of graft duration had immunosuppressant switched to everolimus.

Medical therapy changes after CCTA was more frequent in patients with diabetes ($p=0,043$), but no other association was found with other cardiovascular risk factors, sex, age, previous PA/CVD, ischemic etiology, graft duration, plaque characteristics, calcium score, CAV classification nor previous rejection ($p>0,05$).

Discussion

This real-life single centre experience of CCTA in CAV monitoring in HT patients reinforces the safety of this non-invasive technique, with limited need for additional ischemic or invasive testing.

The potential problem of the negative effect of high resting heart rates on image quality was surpassed using negative chronotropic drugs pre-examination. As CA, CCTA also uses iodine contrast in an already vulnerable population for acute kidney injury and it is fundamental to anticipate who will require contrast nephropathy prophylaxis. The potential lower use of radiation with CCTA is an advantage in this population needing long life monitoring of CAV. A meta-analysis of 13 prospective CCTA studies in 615 HT patients showed mean weighted 94% sensitivity, 92% specificity, 99% negative predictive value, and 67% positive predictive value for detecting stenosis 50% on invasive angiography and the addition of quantitative plaque analysis may also improve sensitivity for CAV detection [5,6].

After the end of our study, new ISHLT guidelines were pre-published in December 2022, giving CCTA an update recommendation as it may be used as a non-invasive alternative to coronary angiography for the detection of CAV in ≥ 2 mm epicardial vessels (class IIa, Level of Evidence (LOE) B). In parallel, also PET was a recommendation class IIa for screening of CAV. On the other side, Stress echocardiogram, MPS and CMR myocardial perfusion reserve have low sensitivity for CAV detection; nevertheless may be useful for prognostication in HT recipients unable to undergo invasive evaluation, CCTA or PET (class IIb, level of evidence B/C) [7].

Similarly to previous studies, our study also supports antibody mediated rejection as a risk factor for the development of CAV – the produced antibodies damage endothelial cells of the coronary arteries of the allograft, triggering an inflammatory response and atherosclerotic plaque formation [8,9].

CCTA adds important information on plaque burden, which may guide therapeutic changes. In fact, after CCTA, therapeutic changes occurred in about one quarter of patients, mainly related to anti-lipidemic optimisation, but also regarding anti-thrombotic drugs and change of immunosuppressants. These changes were more frequent in patients with diabetes, reinforcing the need to intensify anti-lipidemic therapy in this population at high risk of coronary events, despite the possibility of interactions between high-intensity statins and immunosuppressive medication and their side effects. More studies are needed to assess how CCT may guide therapy according to plaque burden.

Limitations

One limitation of this study is the small sample size, which may affect the generalizability and statistical power of our findings. CCTA has limited spatial resolution, not allowing assessment of arteries <2 mm in diameter. We did not perform a direct and simultaneous comparison of CA and CCTA for assessing sensitivity and specificity of CCTA, given that it could not be advantageous for the patient and would increase the risks of exposure to radiation and contrast nephropathy.

Conclusions

In this cohort, CCTA allowed detailed and accurate assessment of CAV with limited need for additional invasive testing. CCTA led to therapeutic changes in about one quarter of patients, and this was more frequent in diabetic patients. More studies are needed to assess how CCT may guide therapy according to plaque burden.

References

1. Lund LH, Edwards LB, Kucheryavaya AY, et al. The registry of the International Society for Heart and Lung Transplantation: thirty-second official adult heart transplantation report–2015; focus theme: early graft failure. *J Heart Lung Transplant* 2015;34:1244-54.
2. Mehra MR, Crespo-Leiro MG, Dipchand A, et al. International Society for Heart and Lung Transplantation working formulation of a standardized nomenclature for cardiac allograft vasculopathy-2010. *J Heart Lung Transplant* 2010;29:717-27.
3. Khush KK, Cherikh WS, Chambers DC, et al. International society for heart and lung transplantation. The International Thoracic Organ Transplant Registry of the International

Society for Heart and Lung Transplantation: thirty-sixth adult heart transplantation report - 2019; focus theme: donor and recipient size match. *J Heart Lung Transplant* 2019;38:1056-66.

4. Costanzo MR, Dipchand A, Starling R, et al. The International Society of Heart and Lung Transplantation guidelines for the care of heart transplant recipients. *J Heart Lung Transplant* 2010;29:914-56.

5. Wever-Pinzon O, Romero J, Kelesidis I, et al. Coronary computed tomography angiography for the detection of cardiac allograft vasculopathy: a meta-analysis of prospective trials. *J Am Coll Cardiol* 2014;63:1992-2004.

6. Miller RJH, Kwiecinski J, Shah KS, et al. Coronary computed tomography-angiography quantitative plaque analysis improves detection of early cardiac allograft vasculopathy: a pilot study. *Am J Transplant* 2019;20:1375-83.

7. Velleca A, Shullo MA, Dhital K, et al. The International Society for Heart and Lung Transplantation (ISHLT) guidelines for the care of heart transplant recipients. *J Heart Lung Transplant* 2023;42:e1-e141.

8. Clerkin KJ, Restaino SW, Zorn E, et al. The effect of timing and graft dysfunction on survival and cardiac allograft vasculopathy in antibody-mediated rejection. *J Heart Lung Transplant* 2016;35:1059-66.

9. Coutance G, Ouldamar S, Rouvier P, et al. Late antibody-mediated rejection after heart transplantation: mortality, graft function, and fulminant cardiac allograft vasculopathy. *J Heart Lung Transplant* 2015;34:1050-7.



Figure 1. Representative coronary computed tomography angiography image of a 56 year-old male heart transplant patient. Anterior descending artery of good calibre, with a non calcified plaque at its medium segment, conditioning a luminal stenosis of up to 25%.

Table 1. Heart transplant patients' characteristics.

Cause of transplantation	
- Dilated cardiomyopathy	16/38 (42.1%)
- Ischemic cardiopathy	12/38 (31.6%)
- Others*	10/18 (26.3%)
Median graft duration, in years	10 (IQR 9)
Median left ventricle ejection fraction, %	62 (IQR 6)
Dyslipidemia	28/38 (73.7%)
Hypertension	24/38 (63.2%)
Diabetes Mellitus	14/38 (36.8%)
Smoking history	10/38 (26.3%)
Obesity	6/38 (15.8%)
Peripheral arterial disease	4/38 (10.5%)
Cerebrovascular disease	4/38 (10.5%)
Medicated with Aspirin or clopidogrel	33/38 (86.8%)
Medicated with Anti-coagulant	4/38 (10.5%)
Medicated with lipid-lowering agent	38/38 (100%)
Statin alone	27/38 (71.1%)
Ezetimibe alone	1/38 (2.6%)
Statin and Ezetimibe	10/38 (26.3%)
Median serum creatinine, mg/dl	1,5 (IQR 0.5)
Median glomerular filtration rate**, ml/min/1.73m ²	56 (IQR 39)
Glomerular filtration rate categories, ml/min/1.73m ²	
Grade 1 (≥ 90 ml/min/1.73m ²)	1/38 (2.6%)
Grade 2 (60-89 ml/min/1.73m ²)	15/38 (39.5%)
Grade 3a (45-59 ml/min/1.73m ²)	9/38 (23.7%)
Grade 3b (30-44 ml/min/1.73m ²)	6/38 (15.8%)
Grade 4 (15-29 ml/min/1.73m ²)	4/38 (10.5%)
Grade 5 (<15 ml/min/1.73m ²)	3/38 (7.9%)

IQRm interquartile range; *other causes, with one or two cases which: valvular disease, myocarditis, cardiotoxicity, Fontan Failure; **glomerular filtration rate, estimated by Modified Modification of Diet in Renal Disease (MDRD) 4-Variable Equation, including sex, age, black race and serum creatinine (mg/dl).

Table 2. Coronary computed tomography angiography characterization – some technical aspects and findings.

Median dose-length product, in mGy*cm	233 (IQR 297)
Median calcium score	17 (IQR 231)
Quality of images	
Excellent	3/32 (9.4%)
Good	21/32 (65.6%)
Moderate	8/32 (25.0%)
ISHLT – CAV classification	
CAV 0	7/32 (21.9%)
CAV 1	24/32 (75.0%)
CAV 2	1/32 (3.1%)
CAV 3	0/32 (0%)
Coronary plaques	
None	7/32 (21.9%)
Exclusively calcified	5/32 (15.6%)
Exclusively non calcified	8/32 (25.0%)
Both calcified and non-calcified	12/32 (37.5%)
Median of affected coronary segments	2 (IQR 3)
Extra-coronary findings	
Cardiac thrombus	4/32 (12.5%)
Pulmonary nodules	2/32 (6.3%)

IQR, interquartile range; IHLTS, International Heart Lung Transplant Society; CAV, cardiac allograft vasculopathy.