

SUPPLEMENTARY MATERIAL

doi: 10.4081/monaldi.2024.3054

Additional yield of transbronchial cryo-node biopsy over endobronchial ultrasound-guided transbronchial needle aspiration for mediastinal lesions at a tertiary care center in India (COLD-FORCEPS-2 study)

Manu Madan,¹ Rohit Kumar,¹ Pranav Ish,¹ Yash Kedia,¹ AJ Mahendran,²
Rajnish Kaushik,¹ Neeraj Kumar Gupta,¹ Nitesh Gupta¹

¹Department of Pulmonary, Critical Care and Sleep Medicine, VMMC and Safdarjung Hospital, Delhi, India; ²SpR Transplant (lung) Medicine, Royal Brompton and Harefield Medicine, London, UK

Correspondence: Nitesh Gupta, Room no. 638, 6th Floor, SSB building, Department of Pulmonary Critical Care and Sleep Medicine, VMMC and Safdarjung Hospital, Delhi, New Delhi - 110029, India.

E-mail-niteshgupta2107@gmail.com

Key words: lymph nodes, mediastinum, endoscopic ultrasound-guided fine needle aspiration/methods, ultrasonography, interventional.

Supplementary Table 1. Comparison of disease-wise additional diagnostic yield of EBUS-TBNA guided mediastinal cryobiopsy over EBUS-TBNA.

	Malignancy	Tuberculosis	Sarcoidosis	Sinus Histiocytosis
Malignancy	-	0.01	0.27	0.001
Tuberculosis	0.01	-	0.058	0.051
Sarcoidosis	0.27	0.058	-	0.001
Sinus histiocytosis	0.001	0.051	0.001	-

Values represent p values of additional yield between two diseases.