



Monaldi Archives for Chest Disease

eISSN 2532-5264

<https://www.monaldi-archives.org/>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community. These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Monaldi Archives for Chest Disease** is, therefore, e-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

All legal disclaimers applicable to the journal apply to this production process as well.

Monaldi Arch Chest Dis 2025 [Online ahead of print]

To cite this Article:

Castro M, Tinoco M, Martins R, et al. **Ghostly intrusion on a frightful Halloween night: a case report of dual valve endocarditis.** *Monaldi Arch Chest Dis* doi: 10.4081/monaldi.2025.3228

 ©The Author(s), 2025
Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.



Ghostly intrusion on a frightful Halloween night: a case report of dual valve endocarditis

Margarida Castro,¹ Mariana Tinoco,¹ Rafael Martins,² Mário Jorge Amorim,²
Luísa Pinheiro,¹ Marina Fernandes,¹ Filipa Cardoso,¹ Filipa Almeida,¹
Paulo Pinho,² João Português,¹ António Lourenço¹

¹Cardiology Department, Local Health Unit of Alto Ave, Guimarães; ²Cardiothoracic Surgery Department, Local Health Unit of São João, Porto, Portugal

Correspondence: Margarida Castro, Cardiology Department, Local Health Unit of Alto Ave, Guimarães, Portugal. Tel.: +351 253 540 330. E-mail: amargaridamcastro@gmail.com

Contributions: MC, MT contributed equally; MC, MT wrote the manuscript with contributions from RM and LP; MJA, MF, FC, FA, PP, JP, AL, reviewed the final version of the manuscript. All authors attest they meet the current ICMJE criteria for authorship, participated in the development of the manuscript and approved its final version.

Conflict of interest: the authors declare no potential conflict of interest.

Ethics approval and consent to participate: no ethical committee approval was required for this case report by the Department, because this article does not contain any studies with human participants or animals.

Patient consent for publication: the patient gave his consent to use his personal data for the publication of this case report and any accompanying images.

Availability of data and materials: all data and materials are available from the corresponding author upon request.

Funding: none.

Abstract

Multivalvular endocarditis (MVE) is an uncommon presentation and mostly involves mitral and aortic valves. Here, we present a case of an MVE with an unusual and bizarre presentation on a Halloween night with a massive degree of valve destruction and right- and left-side involvement requiring emergent surgery.

A 51-year-old male patient with intravenous drug usage presented with anorexia, fever, and dyspnea, rapidly progressing to septic shock with multiorgan dysfunction. Initial blood cultures detected meticillin-sensitive *Staphylococcus aureus*, and antibiotic therapy was started. Transesophageal echocardiography revealed extensive valve destruction of both mitral and tricuspid valves, namely a mitral valve with large vegetation resembling a ghostly figure, causing severe mitral regurgitation. The patient was transferred for emergent mitral and tricuspid surgery.

The particularity of this case, besides the bizarre images, is that, as a drug user, there are specific considerations regarding surgical strategy and options in this scenario that we discussed here.

Key words: infective endocarditis, reconstructive valvular surgery, *Staphylococcus aureus*, embolic risk.

Introduction

Infective endocarditis (IE) continues to be associated with high mortality and may present in a clinically dramatic manner [1]. Multivalvular endocarditis (MVE) is an uncommon presentation and mostly involves mitral and aortic valves [1]. It is associated with higher risk of congestive acute heart failure and septic shock [1]. Vegetations can acquire bizarre shapes with a high degree of valve destruction and often require emergent surgery.

Case Report

We present a case of a 51-year-old-male patient with schizoaffective disorder and cocaine and heroin abuse. He presented to the emergency room on Halloween night due to anorexia, fever and progressive worsening of dyspnoea in the past week. Physical examination showed a systolic murmur and diffuse bilateral crackles on lung auscultation. His clinical status rapidly progressed to septic shock with multiorgan dysfunction and severe dyspnoea with the need for inotropic support and invasive mechanical ventilation (IMV). Initial blood cultures detected Meticillin-Sensitive Staphylococcus aureus (MSSA) and targeted antibiotic therapy was started. Transoesophageal echocardiography revealed a thickened mitral valve with a large, mobile vegetation (maximum 20mm), resembling a ghostly figure (Figure 1 A and B), attached on the posteromedial commissure and interatrial septum with perforation of the posterior leaflet (P2 and P3), causing severe mitral regurgitation (Figure 1D). An abscess in the posteromedial commissure with formation of a pseudoaneurysm was also documented (Figure 1B). Tricuspid valve was also thickened with a vegetation on the septal leaflet, causing moderate tricuspid regurgitation (Figure 1C).

CT scan showed extensive pulmonary (Figure 2A) and splenic (Figure 2B) infarctions.

The patient was transferred to a tertiary surgical center for mitral and tricuspid valve surgery due to extensive valve destruction. He required mitral valve A3 segment, posteromedial commissure, and P3 resection; debridement of a posteroinferior left ventricular free wall abscess (extending to the coronary sinus); and tricuspid anterosseptal commissure, along with anterior and septal leaflet resection.

Reconstructive surgery consisted in closing the left ventricular abscess with autologous pericardium, replacing the excised mitral valve scallops with an autologous pericardial flap, and performing mitral valve annuloplasty with a Physio I annuloplasty ring (Carpentier-Edwards®) (Figure 3). This was followed by replacing the tricuspid excised leaflet segments with an autologous pericardial flap, securing them with two polytetrafluoroethylene (PTFE) cords anchored to the free edge of the neo-anterior and commissural leaflets, and performing annuloplasty with a Contour 3D Ring (Medtronic®).

Transthoracic post operative echocardiographic revealed good biventricular function and normal tricuspid and mitral valve function.

MSSA was isolated from intraoperative valve tissue.

The patient was hospitalized for 44 days with a postoperative period complicated by mediastinitis and nosocomial pneumonia treated conservatively with targeted antibiotic therapy. He was discharged for infectious disease, cardiology and rehabilitation consultations.

Discussion

Half of cases of IE are treated surgically due to important complications [2]. Although this option entails higher operative risk in the acute phase of the disease, it can diminish the risk of heart failure progression, irreversible structural damage and systemic embolism [3].

In this case, the size of vegetation, the embolic events related to both valves' destruction and the refractory shock and heart failure made it imperative to move on to emergent surgery.

In patients with active IV drug abuse, reinfection and mortality rates remain a major challenge [4]. This contributes to the dilemma associated to the best surgical approach as valve replacement poses a heightened risk of recurrence unless absolute abstinence is achieved [4]. Being so, as an IV drug user, there are specific considerations regarding surgical strategy and options. The surgical strategy entails extensive debridement of infected tissues, structural reconstruction, and valve repair if possible [5].

Conclusions

This case reflects the severity and complexity of a bilateral MVE with multiple right- and left-sided septic embolization caused by MSSA in an IV drug user with massive native valve destruction resembling a ghostly figure on a Halloween night.

References

1. Delgado V, Marsan NA, de Waha S, et al. 2023 ESC guidelines for the management of endocarditis. Developed by the task force on the management of endocarditis of the European Society of Cardiology. *Eur Heart J* 2023;44:3948-4042.
2. Lazar HL. Valve surgery for endocarditis in patients who inject drugs: removing them from the society of thoracic surgeons database is only part of the solution. *J Am Heart Assoc* 2021;10:e021153.
3. Zubarevich A, Szczechowicz M, Osswald A, et al. Surgical treatment of infective endocarditis in intravenous drug abusers. *J Cardiothorac Surg* 2021;16:97.
4. Østerdal OB, Salminen P-R, Jordal S, et al. Cardiac surgery for infective endocarditis in patients with intravenous drug use. *Interact CardioVasc Thorac Surg* 2016;22:633-40.
5. Baddour LM, Weimer MB, Wurcel AG, et al. Management of infective endocarditis in people who inject drugs: a scientific statement from the American Heart Association. *Circulation* 2022;146:e187-201.

List of abbreviations

IE	Infective endocarditis
IMV	Invasive mechanical ventilation
IV	Intravenous
MSSA	Meticillin-Sensitive Staphylococcus aureus
MVE	Multivalvular endocarditis
PTFE	Polytetrafluoroethylene




Figure 1. A,B) Transthoracic echocardiogram images showing a thickened mitral valve with a large vegetation resembling a ghostly figure; C) tricuspid valve thickened with a vegetation on the septal leaflet; D) vegetation depending on the posteromedial commissure and interatrial septum with perforation of the posterior leaflet.




Figure 2. Computed tomography scan images showing multiple pulmonary (A) and splenic (B) embolic infarctions.




Figure 3. Reconstructive surgery by replacing the excised mitral valve scallops with an autologous pericardial flap, and performing mitral valve annuloplasty with a Physio I annuloplasty ring (Carpentier-Edwards®). A) Anterior mitral valve leaflet; B) annuloplasty ring; C) autologous pericardial flap; D) posterior mitral valve leaflet.